

Extensor Hi-Fidelity PCR Enzyme Mix

Description: The Extensor Hi-Fidelity PCR Enzyme Mix is a blend of ThermoPrime Taq DNA Polymerase and a proprietary proofreading enzyme. The two enzymes act synergistically, forming a mix that can amplify DNA fragments with double the yields of *Pfu* and at least four times higher fidelity than standard *Taq* DNA polymerase. The system can also generate larger products than either enzyme individually (> 20kb). This makes the enzyme mix ideal for the full-length amplification of sequences for cloning.

Kit Contents:

Vial	Pack Size (cap color)	
	A	B
Extensor Hi-Fidelity PCR Enzyme	20µl (clear)	100µl (clear)
Extensor Hi-Fidelity PCR Buffer 1	1.5ml (red)	5 x 1.5ml (red)
Extensor Hi-Fidelity PCR Buffer 2	1.5ml (green)	5 x 1.5ml (green)
Extensor Hi-Fidelity PCR MgCl ₂	1.5ml (brown)	5 x 1.5ml (brown)
Extensor Hi-Fidelity Primer 1	10µl (clear)	50µl (clear)
Extensor Hi-Fidelity Primer 2	10µl (clear)	50µl (clear)
Extensor Hi-Fidelity Genomic DNA	20µl (clear)	100µl (clear)

The Extensor Hi-Fidelity PCR Enzyme Mix comes supplied with 10X Extensor Buffer 1 (red cap) for PCR < 12kb, 10X Extensor Buffer 2 (green cap) for PCR > 12kb and a vial of 25mM MgCl₂ (brown cap) for any supplementary optimization that may be required. The 10X Buffers both contain 22.5mM MgCl₂.

Storage Conditions: Store at -20°C in a constant temperature freezer for up to 1 year. Avoid freeze thawing. Shipped on ice within the UK and on dry ice for international and within the US.

Protocol:

It is recommended that two master mixes are prepared (on ice). Mix dNTPs, primers and template in one master mix; buffer, enzyme and MgCl₂ in the other. This eliminates the need for a hot start, as well as avoiding any degradation of primers and template through the proofreading activity present in the Extensor Hi-Fidelity PCR Enzyme Mix. The use of wax is not recommended, as it prevents adequate mixing of reaction components, leading to low yields. All reaction tubes should be free of DNase and DNA contamination. Increase enzyme, primer and dNTP concentration for long templates.

Mix together the following:

Master Mix 1	Volume per 25µl reaction
10X Extensor Buffer 1 or 2 (22.5mM MgCl ₂) ¹	2.5µl
Extensor Hi-Fidelity PCR Enzyme Mix (5U/µl) ²	0.125–0.25µl
MgCl ₂ (if required for optimization)	x
Distilled water	to 25µl

Master Mix 2	Volume per 25µl reaction
20mM dNTPs (5mM of each dNTP) ³	1–2.5µl
Template	x
Upstream primer (to final conc. of 0.2µM) ⁴	x
Downstream primer (to final conc. of 0.2µM) ⁴	x
Distilled water	to 25µl

1 - Use Buffer 1 for amplifications up to 12kb. Use Buffer 2 for amplifications longer than 12kb or problematic amplifications of any length.

2 - For long PCR use 0.5µl of enzyme in a 50µl reaction.

3 - For long PCR applications, a final concentration of 350µM–500µM of each dNTP is recommended. For Hi-Fidelity PCR 200–350µM is sufficient.

4 - Primers can be used at 0.4µM for very long extensions.

Mix both master mixes just prior to reaction cycling.

The following points should also be noted:

- Ensure proper mixing of components and always use thin-walled PCR tubes.
- Use a mineral oil overlay unless a heated lid thermocycler is used.
- Touchdown PCR may increase PCR product specificity.
- For best results, use primers of lengths 22–34 nucleotides with annealing temperatures over 60°C.

Templates: For the amplification of large DNA fragments, the quality of the template DNA is very important, as are the denaturation conditions. Keep template DNA denaturation steps as short as possible. Use Extensor Buffer 2 for DNA templates > 12kb and when difficulties are expected or encountered. 250ng human genomic DNA is generally sufficient to provide good PCR results. When using simple templates (such as λ DNA), 1–10ng template DNA should prove sufficient; the number of cycles may be reduced by 5 and Extensor Buffer 1 can be used.

Thermal Cycler Programming: For high fidelity PCR, a standard protocol should be used. For long PCR, modifications have to be made. An example of a long PCR thermal cycling program is given:

Initial denaturation	92–94°C ¹	2 min	1 cycle
Denaturation	92–94°C	10 sec	
Annealing	50–68°C ²	30 sec	10 cycles
Extension	68°C ³	x min ⁴	
Denaturation	94°C	10 sec	
Annealing	50–68°C ²	30 sec	15–20 cycles
Extension (+10 s/cycle)	68°C ³	x min ⁴	
Final extension	68°C	7 min	1 cycle

¹ – When amplifying over 15kb, use a denaturation temperature of 92°C.

² – Annealing temperature dependent on primers.

³ – Always use an extension temperature of 68°C, if possible. Often good results are obtained using a single annealing/extension step at 68°C.

⁴ – Extension times depend on the length of sequence to be amplified (see table below).

Amplicon size (kb)	3	6	10	20	30	40
Extension time (min.)	2	4	8	15	20	30

Ordering Information:	AB-0721/A	Extensor Hi-Fidelity PCR Enzyme Mix	100 units
	AB-0721/B	Extensor Hi-Fidelity PCR Enzyme Mix	500 units

Troubleshooting:

1. *No product detected*
Try reducing the annealing temperature, increasing the concentration or quality of template, concentration of MgCl₂, the number of cycles or improving the purity of primers used.
2. *Spurious bands appearing on electrophoresis gel*
When non-specific products are amplified, try increasing the annealing temperature (up to a maximum of 68°C) or reducing primer concentration, template concentration or cycle number.

For further troubleshooting, see www.abgene.com/troubleshoot.asp or contact Thermo Fisher Scientific (ABgene) TechSupport at abgene.techsupport@thermofisher.com

UK TechSupport, call +44 (0) 1372 840 410

For all other regions, please contact your local Thermo Fisher Scientific (ABgene) office / distributor.

Use of this product is covered by one or more of the following US patents and corresponding patent claims outside the US: 5,079,352, 5,789,224, 5,618,711, 6,127,155 and claims outside the US corresponding to US Patent No. 4,889,818. The purchase of this product includes a limited, non-transferable immunity from suit under the foregoing patent claims for using only this amount of product for the purchaser's own internal research. No right under any other patent claim (such as the patented 5' Nuclease Process claims in US Patents Nos. 5,210,015 and 5,487,972), no right to perform any patented method, and no right to perform commercial services of any kind, including without limitation reporting the results of purchaser's activities for a fee or other commercial consideration, is conveyed expressly, by implication, or by estoppel. This product is for research use only. Diagnostic uses under Roche patents require a separate license from Roche. Further information on purchasing licenses may be obtained by contacting the Director of Licensing, Applied Biosystems, 850 Lincoln Centre Drive, Foster City, California 94404, USA.