

## ABsolute™ Blue QPCR ROX Mix

### Description

ABsolute™ Blue QPCR ROX Mix has been developed to quantify DNA and cDNA\*. With the exception of primers and template, this 2X mix contains all the components required to perform a rapid, sensitive and reproducible QPCR reaction:

- Thermo-Start™ DNA Polymerase, a chemically modified hot-start version of Thermoprime Plus DNA Polymerase, which prevents non-specific amplification during the reaction set-up. **This enzyme requires an activation step at 95°C for 15 minutes.**
- Proprietary reaction buffer which provides highly sensitive, specific and consistent fluorescence readings for real-time and end-point analysis. This buffer has been optimized for MgCl<sub>2</sub> and enhancers to improve amplification across a wide range of templates including plant DNA and GC rich fragments. It contains an inert blue dye to assist in the visualization of the ABsolute Blue QPCR ROX Mix after aliquoting into the reaction well.
- dNTP's, including dTTP to improve reaction sensitivity and efficiency compared to dUTP.
- ROX, passive reference dye for normalization of data.

### Kit Contents

Vial	Pack Size (cap color)	
	A	B
ABsolute Blue QPCR ROX Mix (2X)	2 x 1.25ml (clear)	16 x 1.25ml (clear)

### Cycler & Probe Compatibility

ABsolute™ Blue QPCR ROX Mix is compatible for use with any probe system and QPCR cyclers requiring high ROX dye level, including ABI PRISM® 7000, 7300, 7700, 7900 and 7900HT (including Fast-Block).

\* For RNA template, use Verso 1-Step QRT-PCR ROX Kit (AB-4101)

**INFORMATION****Thermo-Start™ DNA Polymerase**

The enzyme requires an activation step at 95°C for 15 minutes.

Thermo-Start™ has 5' to 3' polymerization and exonuclease activity but lacks 3' to 5' exonuclease activity (proofreading).

**Blue Dye**

This proprietary inert blue dye allows quick and easy visualization of the amount of the mix in the well, minimizing aliquoting errors. It does not interfere with the QPCR reaction and is only available in master mix format.

**ROX Dye**

ROX is an internal passive reference dye used to normalize the fluorescent reporter signal generated in QPCR. The concentration of ROX in the final 1X reaction is 500 nM.

**Storage Conditions**

Store at -20°C until ready for use. Absolute™ Blue QPCR ROX Mix is stable for a minimum of 12 months. The reagents can be stored at 4°C for up to 1 month. Avoid repeated freeze thawing. The ROX dye is light sensitive; exposure should be minimized. Shipped on ice within the UK and on dry ice for international and within the US.

**Additional Info**

- The use of disposable gloves, DNase and RNase free filter tips and plastics is recommended.
- For optimal results, the recommended amplicon length is in the range of 60 to 300 bp.
- As best performance is achieved with dTTP, the Absolute Blue QPCR ROX Mix contains a nucleotide mix with dTTP instead of dUTP.

DIRECTIONS FOR USE

**Tips and Protocol**

Thaw the reagents on ice. Mix and spin down the solutions before use to recover the maximum amount. **Do not vortex the ABsolute Blue QPCR ROX Mix.** Briefly centrifuge to avoid bubbles within the wells, as these will interfere with the fluorescence. Always include a no template control (NTC).

Example of Reaction Mix preparation for a 25 µl final reaction:

		Volume	Final Concentration
Reaction Mix	ABsolute Blue QPCR ROX Mix (2X)	12.5 µl	1X
	Forward primer (10 µM) <sup>a</sup>	1 µl	400 nM
	Reverse primer (10 µM) <sup>a</sup>	1 µl	400 nM
	Probe	Variable	100 - 250 nM
	Water (PCR grade) <sup>b</sup>	Variable	
	Template (DNA or cDNA) <sup>c</sup>	1 - 5 µl	<250 ng/reaction
Total volume		25 µl	

Example of a QPCR thermal cycling program:

	Temp.	Time	Number of cycle
Enzyme activation	95°C	<b>15 min</b>	1 cycle
Denaturation	95°C	15 sec	40 cycles
Annealing/Extension <sup>d</sup>	60°C	60 sec	

**Notes**

- a – For optimization, a primer titration should be performed from 100 nM to 500 nM final concentration. Scale up or down the volume and concentration as appropriate.
- b – The volume of the total reaction should be completed up to 25 µl with water.
- c – The volume of template to add to the QPCR reaction can be adjusted as required. For standard templates only 1 µl should be added to reduce carryover of PCR inhibitors. This volume can be increased up to 5 µl for low copy number templates.
- d – Separate annealing (50–60°C for 30 sec) and extension steps (72°C for 30 sec) may be necessary with some probe systems (e.g. Molecular Beacons), as the optimal temperature for detecting fluorescence may be different.

## Quality control

ABsolute Blue QPCR ROX Mix is tested functionally using QPCR. The product must demonstrate linearity of amplification over a specified serial dilution of human genomic DNA.

## Ordering Information

AB-4138/A	ABsolute™ Blue QPCR ROX Mix	200 x 25 µl rxns
AB-4138/B	ABsolute™ Blue QPCR ROX Mix	1,600 x 25 µl rxns
AB-4139/A	ABsolute™ Blue QPCR ROX Mix	400 x 25 µl rxns
AB-4139/B	ABsolute™ Blue QPCR ROX Mix	4,000 x 25 µl rxns

## Related Products

Cat. No.	Description	Quantity
AB-1100/W	Thermo-Fast™ 96 PCR Detection Plate, white *	25 plates
AB-1400/W	Thermo-Fast™ 96 PCR Detection Plate Mark II, white *	25 plates
AB-1170	ABsolute™ QPCR Seal (adhesive seal)	50 sheets
AB-0812	Clear Seal Diamond (heat seal)	100 sheets
AB-0866	Ultra Clear Cap Strips (8 caps)	120 strips

\* For Cycler compatibility and other color choices, see our latest catalogue or visit [www.abgene.com](http://www.abgene.com)

## Troubleshooting

For troubleshooting, see [www.abgene.com/troubleshoot.asp](http://www.abgene.com/troubleshoot.asp) or contact Thermo Fisher Scientific (ABgene) TechSupport at [abgene.techsupport@thermofisher.com](mailto:abgene.techsupport@thermofisher.com)

UK TechSupport, call +44 (0) 1372 840 410

**For all other regions, please contact your local Thermo Fisher Scientific (ABgene) office / distributor.**

A license to perform the patented 5' Nuclease Process for research is obtained by the purchase of (i) both Authorized 5' Nuclease Core Kit and Licensed Probe, (ii) a Licensed 5' Nuclease Kit, or (iii) license rights from Applied Biosystems.

This product is an Authorized 5' Nuclease Core Kit. Use of this product is covered by one or more of the following US patents and corresponding patent claims outside the US: 5,079,352, 5,789,224, 5,618,711, 6,127,155, 5,677,152, 5,773,258, 5,407,800, 5,322,770, 5,310,652, 5,210,015, 5,487,972, and claims outside the US corresponding to US Patent No. 4,889,818. The purchase of this product includes a limited, non-transferable immunity from suit under the foregoing patent claims for using only this amount of product for the purchaser's own internal research. Separate purchase of a Licensed Probe would convey rights under the applicable claims of US Patents Nos. 5,538,848, 5,723,591, 5,876,930, 6,030,787, 6,258,569, 5,804,375 (claims 1-12 only), and 6,214,979, and corresponding claims outside the United States. No right under any other patent claim and no right to perform commercial services of any kind, including without limitation reporting the results of purchaser's activities for a fee or other commercial consideration, is conveyed expressly, by implication, or by estoppel. This product is for research use only. Diagnostic uses under Roche patents require a separate license from Roche. Further information on purchasing licenses may be obtained from the Director of Licensing, Applied Biosystems, 850 Lincoln Centre Drive, Foster City, California 94404, USA.

**For Research Purposes Only**

4